

# From Classical Gravity to Quantum Amplitudes (lecture 2b)

**Thibault Damour**

Institut des Hautes Etudes Scientifiques



**Institut de Physique Théorique *and*  
Institut des Hautes Études Scientifiques**

***Cours de Physique Théorique***

**IPhT, Fridays 5, 12 October 2018 (10:00 to 12:15),**

**and Friday 19 October (10:00 to 12:15, and 14:15 to 16:30)**

# PN, EOB, NR, PHENOMD

PN accuracy loss during late inspiral


-> **necessity to develop an improved description**

Dimensionless « quality factor » of GW phase  $Q_\omega = f^2 \frac{d^2 \psi(f)}{df^2} \approx \frac{\omega^2}{\dot{\omega}}$

**difference**  
from LO

$$Q_\omega - Q_\omega^N$$

**$v=1/2$  at merger !**



# Analytical GW Templates for BBH Coalescences ?

PN corrections to Einstein's quadrupole frequency « chirping »  
 from PN-improved balance equation  $dE(f)/dt = - F(f)$

$$\frac{d\phi}{d \ln f} = \frac{\omega^2}{d\omega/dt} = Q_\omega^N \hat{Q}_\omega$$

$$Q_\omega^N = \frac{5c^5}{48\nu v^5}; \hat{Q}_\omega = 1 + c_2 \left(\frac{v}{c}\right)^2 + c_3 \left(\frac{v}{c}\right)^3 + \dots$$

$$\frac{v}{c} = \left( \frac{\pi G(m_1 + m_2) f}{c^3} \right)^{\frac{1}{3}}$$


$$\nu = \frac{m_1 m_2}{(m_1 + m_2)^2}$$

**Cutler et al. '93:**

« slow convergence of PN »

**Brady-Creighton-Thorne'98:**


« inability of current computational techniques to evolve a BBH through its last ~10 orbits of inspiral » and to compute the merger




**Damour-Iyer-Sathyaprakash'98:**

use **resummation** methods for E and F

**Buonanno-Damour '99-00:**  
 novel, resummed approach:  
**Effective-One-Body**  
**analytical formalism**







# Effective One Body (EOB) Method

Buonanno-Damour 1999, 2000; Damour-Jaranowski-Schaefer 2000; Damour 2001 ;Damour-Iyer-Nagar 2008

Resummation of perturbative PN results  $\longrightarrow$  description of the coalescence  
 + addition of ringdown (Vishveshwara 70, Davis-Ruffini-Tiomno 1972)


Buonanno-Damour 2000



Predictions as early as 2000 :  
 continued transition, non adiabaticity, first complete waveform, final spin (OK within 10%), final mass

# First complete waveforms for BBH coalescences: analytical EOB



Non-spinning BHs  
Buonanno-Damour 2000



Spinning BHs  
Buonanno-Chen-Damour

Nov 2005:

« to show the  
promise  
of a purely  
analytical  
EOB-based  
approach »



# EOB THEORY + EOB[NR] + EOB[SF] DEVELOPMENTS

Buonanno, Damour 99 (2 PN Hamiltonian)  
Buonanno, Damour 00 (Rad.Reac. full waveform)  
Damour, Jaranowski, Schäfer 00 (3 PN Hamiltonian)  
Damour 01, (spinning bodies)  
Buonanno, Chen, Damour 05,  
Damour-Jaranowski, Schäfer 08, Barausse, Buonanno, 10, Nagar 11,  
Balmelli-Jetzer 12, Taracchini et al 12,14, Damour, Nagar 14 (factorized waveform)  
Damour, Nagar 07,  
Damour, Iyer, Nagar 08,  
Pan et al. 11  
Damour, Nagar 10 (BNS tidal effects)  
Bini-Damour-Faye 12  
Bini, Damour 13, Damour, Jaranowski, Schäfer 15 (4 PN Hamiltonian)

## EOB vs NR and EOB[NR]

Buonanno, Cook, Pretorius 07,  
Buonanno, Pan, Taracchini 08-  
Damour-Nagar 08-

Reduced Order Model version (Pürrer 2014, 2016) of EOB[NR] (Taracchini et al 2014)

Phenomenological model (Ajith et al 2007, Hannam et al 2014, Husa et al 2016, Kahn et al 2016) of FFT of hybrids EOB + NR


## EOB vs SF and EOB[SF]

Damour 09  
Barack-Sago-Damour 10  
Barausse-Buonanno-LeTiec 12  
Akçay-Barack-Damour-Sago 12  
Bini-Damour 13-16  
LeTiec 15  
Bini-Damour-Geralico 16  
Hopper-Kavanagh-Ottewill 16  
Akçay-vandeMeent 16

## EOB vs PM

Damour 16

**EOB: resumming the dynamics of a two-body system  $(m_1, m_2, S_1, S_2)$  in terms of the dynamics of a particle of mass  $\mu$  and spin  $S^*$  moving in some effective metric  $g(M, S)$**




**Effective metric for non-spinning bodies: a nu-deformation of Schwarzschild**

$$M = m_1 + m_2 \quad \mu = \frac{m_1 m_2}{m_1 + m_2} \quad \nu = \frac{\mu}{M} = \frac{m_1 m_2}{(m_1 + m_2)^2}$$



$$ds_{\text{eff}}^2 = -A(r; \nu) dt^2 + B(r; \nu) dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\varphi^2)$$

# Real dynamics versus Effective dynamics

Real dynamics



Effective dynamics



$$S = - \int \mu ds + \dots$$

$$H = H_0 + \left( G H_1 + \frac{G^2}{c^2} H_2 + \frac{G^3}{c^4} H_3 + \frac{G^4}{c^6} H_4 \right) \left( 1 + \frac{1}{c^2} + \dots \right)$$

Effective metric for non-spinning bodies: a nu-deformation of Schwarzschild


$$\nu = \frac{m_1 m_2}{(m_1 + m_2)^2}$$

$$ds_{\text{eff}}^2 = -A(r; \nu) dt^2 + B(r; \nu) dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\varphi^2)$$

# TWO-BODY/EOB "CORRESPONDENCE":

THINK QUANTUM-MECHANICALLY (J.A. WHEELER)


Real 2-body system  
(in the c.o.m. frame)  
 $(m_1, m_2)$



An effective particle  
in some effective metric

$$\mu = \frac{m_1 m_2}{m_1 + m_2}$$

$$g_{\mu\nu}^{\text{eff}}$$



$$\mu^2 + g_{\text{eff}}^{\mu\nu} \frac{\partial S_{\text{eff}}}{\partial x^\mu} \frac{\partial S_{\text{eff}}}{\partial x^\nu} + \mathcal{O}(p^4) = 0$$

Bohr-Sommerfeld's  
Quantization Conditions  
(action-angle variables &  
Delaunay Hamiltonian)

$$J = l\hbar = \frac{1}{2\pi} \oint p_\varphi d\varphi$$

$$N = n\hbar = I_r + J$$

$$I_r = \frac{1}{2\pi} \oint p_r dr$$

$$H^{\text{classical}}(q, p) \longrightarrow H^{\text{classical}}(I_a) \longrightarrow E^{\text{quantum}}(I_a = n_a h) = f^{-1}[\mathcal{E}_{\text{eff}}^{\text{quantum}}(I_a^{\text{eff}} = n_a h)]$$

## 2-body Taylor-expanded N + 1PN + 2PN Hamiltonian

$$H_N(\mathbf{x}_a, \mathbf{p}_a) = \frac{\mathbf{p}_1^2}{2m_1} - \frac{1}{2} \frac{Gm_1m_2}{r_{12}} + (1 \leftrightarrow 2)$$

$$c^2 H_{1PN}(\mathbf{x}_a, \mathbf{p}_a) = -\frac{1}{8} \frac{(\mathbf{p}_1^2)^2}{m_1^3} + \frac{1}{8} \frac{Gm_1m_2}{r_{12}} \left( -12 \frac{\mathbf{p}_1^2}{m_1^2} + 14 \frac{(\mathbf{p}_1 \cdot \mathbf{p}_2)}{m_1m_2} + 2 \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)}{m_1m_2} \right) \\ + \frac{1}{4} \frac{Gm_1m_2}{r_{12}} \frac{G(m_1 + m_2)}{r_{12}} + (1 \leftrightarrow 2),$$

$$c^4 H_{2PN}(\mathbf{x}_a, \mathbf{p}_a) = \frac{1}{16} \frac{(\mathbf{p}_1^2)^3}{m_1^5} + \frac{1}{8} \frac{Gm_1m_2}{r_{12}} \left( 5 \frac{(\mathbf{p}_1^2)^2}{m_1^4} - \frac{11}{2} \frac{\mathbf{p}_1^2 \mathbf{p}_2^2}{m_1^2 m_2^2} - \frac{(\mathbf{p}_1 \cdot \mathbf{p}_2)^2}{m_1^2 m_2^2} + 5 \frac{\mathbf{p}_1^2 (\mathbf{n}_{12} \cdot \mathbf{p}_2)^2}{m_1^2 m_2^2} \right. \\ \left. - 6 \frac{(\mathbf{p}_1 \cdot \mathbf{p}_2)(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)}{m_1^2 m_2^2} - \frac{3}{2} \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)^2 (\mathbf{n}_{12} \cdot \mathbf{p}_2)^2}{m_1^2 m_2^2} \right) \\ + \frac{1}{4} \frac{G^2 m_1 m_2}{r_{12}^2} \left( m_2 \left( 10 \frac{\mathbf{p}_1^2}{m_1^2} + 19 \frac{\mathbf{p}_2^2}{m_2^2} \right) - \frac{1}{2} (m_1 + m_2) \frac{27(\mathbf{p}_1 \cdot \mathbf{p}_2) + 6(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)}{m_1 m_2} \right) \\ - \frac{1}{8} \frac{Gm_1m_2}{r_{12}} \frac{G^2(m_1^2 + 5m_1m_2 + m_2^2)}{r_{12}^2} + (1 \leftrightarrow 2),$$

# Technical aspects of EOB (1)

1. Go to the c.m. frame ( $p_1 = -p_2 = p$ )

2. Compute both the real and the effective Delaunay Hamiltonian, i.e.  $H_{\text{real}}(I_{\text{real}})$  and  $H_{\text{eff}}(I_{\text{eff}})$ , functions of the action integrals  $I_a$ , and look for an energy map  $f(E)$ , such that:

$$f[H_{\text{real}}(I_a)] = H_{\text{eff}}(I_a)$$

3. Alternatively, look for a canonical transformation  $G(q, p')$ , and an energy map  $f(E)$ , such that:

$$q'^i = q^i + \frac{\partial G(q, p')}{\partial p'_i}, \quad p_i = p'_i + \frac{\partial G(q, p')}{\partial q^i}$$

$$\boxed{\frac{H_{\text{eff}}^{g_{\mu\nu}}(\mathbf{q}', \mathbf{p}')}{\mu c^2} = \left[ f \left( \frac{H_{\text{real}}(\mathbf{q}, \mathbf{p})}{\mu c^2} \right) \right]_{(\mathbf{q}, \mathbf{p}) \xrightarrow{G} (\mathbf{q}', \mathbf{p}')},}$$

4. Parametrizing a general energy-map:

$$f \left( \frac{E_{\text{real}}}{\mu c^2} \right) = 1 + \frac{E_{\text{real}}^{\text{binding}}}{\mu c^2} \left( 1 + \alpha_1 \frac{E_{\text{real}}^{\text{binding}}}{\mu c^2} + \alpha_2 \left( \frac{E_{\text{real}}^{\text{binding}}}{\mu c^2} \right)^2 + \alpha_3 \left( \frac{E_{\text{real}}^{\text{binding}}}{\mu c^2} \right)^3 + \dots \right).$$

# Computing radial integrals à la Sommerfeld (Damour-Schaefer'88)

$$I_r(E, J) = \frac{1}{2\pi} \oint p_r(E, J, r) dr$$


$$I_\varphi = \frac{1}{2\pi} \oint p_\varphi d\varphi = p_\varphi = J$$

$$I(A, B, C, D_1, D_2, D_3) = \frac{2}{2\pi} \int_{r_{\min}}^{r_{\max}} dr \left( A + \frac{2B}{r} + \frac{C}{r^2} + \frac{D_1}{r^3} + \frac{D_2}{r^4} + \frac{D_3}{r^5} \right)^{\frac{1}{2}}$$

$$(3.9) \quad I(A, B, C, D_1, D_2, D_3) = \frac{B}{\sqrt{-A}} -$$

$$-\sqrt{-C} \left\{ 1 - \frac{1}{2} \frac{B}{C^2} \left[ D_1 - \frac{3}{2} \frac{D_2 B}{C} + \frac{15}{8} \frac{D_1^2 B}{C^2} \right] - \right.$$

$$\left. - \frac{1}{4} \frac{A}{C^2} \left[ D_2 - \frac{3}{4} \frac{D_1^2}{C} \right] + \frac{3}{4} \frac{B}{C^3} \left[ A - \frac{5}{3} \frac{B^2}{C} \right] D_3 \right\} + O(D_1^3 + D_2^2 + D_3^2 + D_1^2 D_2 + \dots).$$



$$\mathcal{E}^R(\mathcal{N}, \mathcal{J}) = Mc^2 - \frac{1}{2} \frac{\mu \alpha^2}{\mathcal{N}^2} \left[ 1 + \frac{\alpha^2}{c^2} \left( \frac{6}{\mathcal{N}\mathcal{J}} - \frac{1}{4} \frac{15-\nu}{\mathcal{N}^2} \right) \right.$$

$$\alpha \equiv \mu GM = Gm_1 m_2 \quad \left. + \frac{\alpha^4}{c^4} \left( \frac{5}{2} \frac{7-2\nu}{\mathcal{N}\mathcal{J}^3} + \frac{27}{\mathcal{N}^2 \mathcal{J}^2} - \frac{3}{2} \frac{35-4\nu}{\mathcal{N}^3 \mathcal{J}} \right) \right]$$

$$N = I_r + I_\varphi = I_r + J \quad \left. + \frac{1}{8} \frac{145-15\nu+\nu^2}{\mathcal{N}^4} \right], \quad (3.10)$$

# Technical aspects of EOB (2)

the coefficients of the spherically-symmetric effective metric,

$$g_{\mu\nu}^{\text{eff}} dx^\mu dx^\nu = -A(R; \nu) c^2 dT^2 + B(R; \nu) dR^2 + R^2(d\theta^2 + \sin^2 \theta d\varphi^2).$$

are looked for as usual-type PN expansions:

$$A(R; \nu) = 1 + \tilde{a}_1 \frac{GM}{c^2 R} + \tilde{a}_2 \left( \frac{GM}{c^2 R} \right)^2 + \tilde{a}_3 \left( \frac{GM}{c^2 R} \right)^3 + \tilde{a}_4 \left( \frac{GM}{c^2 R} \right)^4 + \dots;$$

$$B(R; \nu) = 1 + \tilde{b}_1 \frac{GM}{c^2 R} + \tilde{b}_2 \left( \frac{GM}{c^2 R} \right)^2 + \tilde{b}_3 \left( \frac{GM}{c^2 R} \right)^3 + \dots,$$

System of algebraic eqs for unknown coefficients:  $a_n, b_n, \alpha_n$ ; impose  $b_1=2$ :

-> **unique solution at 2PN** (Buonanno-Damour'99)

$$\alpha_1 = \frac{\nu}{2}, \quad \alpha_2 = 0.$$

->

$$\frac{\mathcal{E}_0}{m_0 c^2} \equiv \frac{\mathcal{E}_{\text{real}}^2 - m_1^2 c^4 - m_2^2 c^4}{2m_1 m_2 c^4}.$$

$$\hat{a}_2 = 0, \quad \hat{a}_3 = 2\nu, \quad \hat{b}_2 = 4 - 6\nu.$$

-> **2PN effective metric**

# EOB results at 1PN and 2PN

**Theorem 1:** The Lorentz-Droste-Einstein-Infeld-Hoffmann 1PN dynamics, considered in the center-of-mass frame, is mapped (at 1PN accuracy) onto the geodesic motion of a particle of mass  $\mu = m_1 m_2 / (m_1 + m_2)$  in a Schwarzschild background of mass  $M = m_1 + m_2$ , modulo the very simple (but non trivial) energy map

$$\frac{\mathcal{E}_{\text{eff}}}{\mu c^2} = \frac{(\mathcal{E}_{\text{real}}^{\text{tot}})^2 - m_1^2 c^4 - m_2^2 c^4}{2 m_1 m_2 c^4} \quad (\text{at the 1PN, 2PN, 3PN, and 4PN levels}).$$

**Theorem 2:** The full 2PN dynamics (whose general-frame Hamiltonian contains thirteen 2PN-level independent terms besides the five 1PN-level ones), when considered in the center-of-mass frame, is mapped (at 2PN accuracy) onto the geodesic motion of a particle of mass  $\mu$  in the following simple  $\nu$ -deformation of the Schwarzschild metric of mass  $M$ ,

$$(ds_{\text{eff}}^2)^{2\text{PN}} = - \left( 1 - 2 \frac{GM}{c^2 R} + 2\nu \left( \frac{GM}{c^2 R} \right)^3 \right) c^2 dt^2 + \frac{1 - 6\nu \left( \frac{GM}{c^2 R} \right)^2}{1 - 2 \frac{GM}{c^2 R}} dR^2 + R^2 (d\theta^2 + \sin^2 \theta d\varphi^2), \quad (65)$$

modulo the same energy map <sup>(energy map)</sup> (54) that appeared at the 1PN level.

## 2-body Taylor-expanded 3PN Hamiltonian [JS 98, DJS 01]

$$\begin{aligned}
 c^6 H_{3\text{PN}}(\mathbf{x}_a, \mathbf{p}_a) = & -\frac{5}{128} \frac{(\mathbf{p}_1^2)^4}{m_1^7} + \frac{1}{32} \frac{Gm_1 m_2}{r_{12}} \left( -14 \frac{(\mathbf{p}_1^2)^3}{m_1^6} + 4 \frac{((\mathbf{p}_1 \cdot \mathbf{p}_2)^2 + 4\mathbf{p}_1^2 \mathbf{p}_2^2) \mathbf{p}_1^2}{m_1^4 m_2^2} + 6 \frac{\mathbf{p}_1^2 (\mathbf{n}_{12} \cdot \mathbf{p}_1)^2 (\mathbf{n}_{12} \cdot \mathbf{p}_2)^2}{m_1^4 m_2^2} \right. \\
 & - 10 \frac{(\mathbf{p}_1^2 (\mathbf{n}_{12} \cdot \mathbf{p}_2)^2 + \mathbf{p}_2^2 (\mathbf{n}_{12} \cdot \mathbf{p}_1)^2) \mathbf{p}_1^2}{m_1^4 m_2^2} + 24 \frac{\mathbf{p}_1^2 (\mathbf{p}_1 \cdot \mathbf{p}_2) (\mathbf{n}_{12} \cdot \mathbf{p}_1) (\mathbf{n}_{12} \cdot \mathbf{p}_2)}{m_1^4 m_2^2} \\
 & + 2 \frac{\mathbf{p}_1^2 (\mathbf{p}_1 \cdot \mathbf{p}_2) (\mathbf{n}_{12} \cdot \mathbf{p}_2)^2}{m_1^3 m_2^3} + \frac{(7\mathbf{p}_1^2 \mathbf{p}_2^2 - 10(\mathbf{p}_1 \cdot \mathbf{p}_2)^2) (\mathbf{n}_{12} \cdot \mathbf{p}_1) (\mathbf{n}_{12} \cdot \mathbf{p}_2)}{m_1^3 m_2^3} \\
 & + \frac{(\mathbf{p}_1^2 \mathbf{p}_2^2 - 2(\mathbf{p}_1 \cdot \mathbf{p}_2)^2) (\mathbf{p}_1 \cdot \mathbf{p}_2)}{m_1^3 m_2^3} + 15 \frac{(\mathbf{p}_1 \cdot \mathbf{p}_2) (\mathbf{n}_{12} \cdot \mathbf{p}_1)^2 (\mathbf{n}_{12} \cdot \mathbf{p}_2)^2}{m_1^3 m_2^3} \\
 & - 18 \frac{\mathbf{p}_1^2 (\mathbf{n}_{12} \cdot \mathbf{p}_1) (\mathbf{n}_{12} \cdot \mathbf{p}_2)^3}{m_1^3 m_2^3} + 5 \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)^3 (\mathbf{n}_{12} \cdot \mathbf{p}_2)^3}{m_1^3 m_2^3} \left. \right) + \frac{G^2 m_1 m_2}{r_{12}^2} \left( \frac{1}{16} (m_1 - 27m_2) \frac{(\mathbf{p}_1^2)^2}{m_1^4} \right. \\
 & - \frac{115}{16} m_1 \frac{\mathbf{p}_1^2 (\mathbf{p}_1 \cdot \mathbf{p}_2)}{m_1^3 m_2} - \frac{1}{48} m_2 \frac{25(\mathbf{p}_1 \cdot \mathbf{p}_2)^2 + 371\mathbf{p}_1^2 \mathbf{p}_2^2}{m_1^2 m_2^2} + \frac{17}{16} \frac{\mathbf{p}_1^2 (\mathbf{n}_{12} \cdot \mathbf{p}_1)^2}{m_1^3} + \frac{5}{12} \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)^4}{m_1^3} \\
 & - \frac{1}{8} m_1 \frac{(15\mathbf{p}_1^2 (\mathbf{n}_{12} \cdot \mathbf{p}_2) + 11(\mathbf{p}_1 \cdot \mathbf{p}_2) (\mathbf{n}_{12} \cdot \mathbf{p}_1)) (\mathbf{n}_{12} \cdot \mathbf{p}_1)}{m_1^3 m_2} - \frac{3}{2} m_1 \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)^3 (\mathbf{n}_{12} \cdot \mathbf{p}_2)}{m_1^3 m_2} \\
 & + \frac{125}{12} m_2 \frac{(\mathbf{p}_1 \cdot \mathbf{p}_2) (\mathbf{n}_{12} \cdot \mathbf{p}_1) (\mathbf{n}_{12} \cdot \mathbf{p}_2)}{m_1^2 m_2^2} + \frac{10}{3} m_2 \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)^2 (\mathbf{n}_{12} \cdot \mathbf{p}_2)^2}{m_1^2 m_2^2} \\
 & - \frac{1}{48} (220m_1 + 193m_2) \frac{\mathbf{p}_1^2 (\mathbf{n}_{12} \cdot \mathbf{p}_2)^2}{m_1^2 m_2^2} \left. \right) + \frac{G^3 m_1 m_2}{r_{12}^3} \left( -\frac{1}{48} \left( 425m_1^2 + \left( 473 - \frac{3}{4} \pi^2 \right) m_1 m_2 + 150m_2^2 \right) \frac{\mathbf{p}_1^2}{m_1^2} \right. \\
 & + \frac{1}{16} \left( 77(m_1^2 + m_2^2) + \left( 143 - \frac{1}{4} \pi^2 \right) m_1 m_2 \right) \frac{(\mathbf{p}_1 \cdot \mathbf{p}_2)}{m_1 m_2} + \frac{1}{16} \left( 20m_1^2 - \left( 43 + \frac{3}{4} \pi^2 \right) m_1 m_2 \right) \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)^2}{m_1^2} \\
 & + \frac{1}{16} \left( 21(m_1^2 + m_2^2) + \left( 119 + \frac{3}{4} \pi^2 \right) m_1 m_2 \right) \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1) (\mathbf{n}_{12} \cdot \mathbf{p}_2)}{m_1 m_2} \left. \right) \\
 & + \frac{1}{8} \frac{G^4 m_1 m_2^3}{r_{12}^4} \left( \left( \frac{227}{3} - \frac{21}{4} \pi^2 \right) m_1 - m_2 \right) + (1 \leftrightarrow 2).
 \end{aligned}$$

## EOB results at 3PN (DJS'00)

Less eqs than unknowns ? Need to introduce higher-in-momenta contributions

$$S = -\mu \int ds_{\text{eff}} [1 + A_{\mu\nu\kappa\lambda}(x) u^\mu u^\nu u^\kappa u^\lambda + \dots].$$

post-geodesic effective mass-shell condition

$$g_{\text{eff}}^{\mu\nu} P'_\mu P'_\nu + \mu^2 c^2 + Q(P'_\mu) = 0,$$

at 3PN

$$Q(P'_\mu) = \frac{1}{c^6} \frac{1}{\mu^2} \left( \frac{GM}{R'} \right)^2 [z_1 \mathbf{P}'^4 + z_2 \mathbf{P}'^2 (\mathbf{n}' \cdot \mathbf{P}')^2 + z_3 (\mathbf{n}' \cdot \mathbf{P}')^4]$$

$$8 z_1 + 4 z_2 + 3 z_3 = 6\nu(4 - 3\nu).$$

DJS gauge:  $z_1=z_2=0$  to have only  $p^4$  terms

# 3PN EOB

**Theorem 3:** The full 3PN dynamics (whose general-frame Hamiltonian contains  $\sim 40$  3PN-level terms), when considered in the center-of-mass frame, is mapped (at 3PN accuracy), via the simple energy map (54) that appeared at 1PN, onto the motion of a particle of mass  $\mu$  submitted to the mass-shell condition (38) where  $g_{\mu\nu}^{\text{eff}}$  is given by Eq. , with

$$A^{3\text{PN}}(u) = 1 - 2u + 2\nu u^3 + \left( \frac{94}{3} - \frac{41}{32} \pi^2 \right) \nu u^4, \quad (71)$$

$$\bar{D}^{3\text{PN}}(u) = 1 + 6\nu u^2 + (52\nu - 6\nu^2) u^3, \quad (72)$$

and where

$$\hat{Q}^{3\text{PN}} \equiv \frac{Q}{\mu^2 c^2} = (8\nu - 6\nu^2) u^2 \frac{p_r^4}{c^4}. \quad (73)$$

Here  $\bar{D} \equiv (AB)^{-1}$ , we used the scaled momentum  $p_r \equiv P_R^{\text{EOB}}/\mu$  (with dimension  $[p] = [\text{velocity}]$ ) so that  $p/c$  is dimensionless, and we introduced the convenient dimensionless EOB variable

$$u \equiv \frac{GM}{c^2 R_{\text{EOB}}}. \quad (74)$$

# 2-body Taylor-expanded 4PN Hamiltonian [DJS, 2014]

$${}^8H_{4PN}^{\text{local}}(\mathbf{x}_a, \mathbf{p}_a) = \frac{7(\mathbf{p}_1^2)^5}{256m^7} + \frac{Gm_1m_2}{r_{12}} H_{40}(\mathbf{x}_a, \mathbf{p}_a) + \frac{G^2m_1m_2}{r_{12}^2} m_1 H_{40}(\mathbf{x}_a, \mathbf{p}_a) \\ + \frac{G^2m_1m_2}{r_{12}^2} (m_1^2 H_{44}(\mathbf{x}_a, \mathbf{p}_a) + m_1m_2 H_{442}(\mathbf{x}_a, \mathbf{p}_a)) \\ + \frac{G^4m_1m_2}{r_{12}^4} (m_1^2 H_{12}(\mathbf{x}_a, \mathbf{p}_a) + m_1^2 m_2 H_{122}(\mathbf{x}_a, \mathbf{p}_a)) \\ + \frac{G^5m_1m_2}{r_{12}^5} H_{40}(\mathbf{x}_a, \mathbf{p}_a) + (1 \leftrightarrow 2), \quad (\text{A3})$$

$$H_{40}(\mathbf{x}_a, \mathbf{p}_a) = \frac{45(\mathbf{p}_1^2)^4}{128m_1^4} - \frac{5(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)^3}{64m_1^3m_2^3} + \frac{15(\mathbf{n}_{12} \cdot \mathbf{p}_2)^2(\mathbf{p}_1^2)^3}{64m_1^3m_2^3} - \frac{9(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)^2(\mathbf{p}_1 \cdot \mathbf{p}_2)}{16m_1^2m_2^2} \\ + \frac{3(\mathbf{p}_1^2)^2(\mathbf{p}_1 \cdot \mathbf{p}_2)^2}{32m_1^2m_2^2} + \frac{15(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)^2(\mathbf{p}_2^2)}{64m_1^2m_2^2} - \frac{2(\mathbf{p}_1^2)(\mathbf{p}_2^2)^2}{64m_1^2m_2^2} + \frac{25(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)}{256m_1^2m_2^2} \\ + \frac{25(\mathbf{n}_{12} \cdot \mathbf{p}_1)^2(\mathbf{n}_{12} \cdot \mathbf{p}_2)^2(\mathbf{p}_1^2)}{128m_1^2m_2^2} + \frac{23(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{256m_1^2m_2^2} - \frac{85(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1 \cdot \mathbf{p}_2)}{256m_1^2m_2^2} \\ + \frac{45(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)(\mathbf{p}_1 \cdot \mathbf{p}_2)}{128m_1^2m_2^2} - \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_2)^2(\mathbf{p}_1^2)(\mathbf{p}_1 \cdot \mathbf{p}_2)}{256m_1^2m_2^2} + \frac{25(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1 \cdot \mathbf{p}_2)^2}{64m_1^2m_2^2} \\ + \frac{7(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1 \cdot \mathbf{p}_2)^2}{64m_1^2m_2^2} - \frac{3(\mathbf{n}_{12} \cdot \mathbf{p}_1)^2(\mathbf{p}_1 \cdot \mathbf{p}_2)^2}{64m_1^2m_2^2} - \frac{3(\mathbf{p}_1^2)(\mathbf{p}_1 \cdot \mathbf{p}_2)^2}{64m_1^2m_2^2} + \frac{55(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{256m_1^2m_2^2} \\ + \frac{7(\mathbf{n}_{12} \cdot \mathbf{p}_1)^2(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{128m_1^2m_2^2} - \frac{22(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{256m_1^2m_2^2} - \frac{23(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{p}_1 \cdot \mathbf{p}_2)(\mathbf{p}_2^2)}{256m_1^2m_2^2} \\ + \frac{7(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{p}_1 \cdot \mathbf{p}_2)(\mathbf{p}_2^2)}{128m_1^2m_2^2} - \frac{7(\mathbf{p}_1^2)(\mathbf{p}_1 \cdot \mathbf{p}_2)(\mathbf{p}_2^2)}{256m_1^2m_2^2} - \frac{5(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{64m_1^2m_2^2} + \frac{7(\mathbf{n}_{12} \cdot \mathbf{p}_2)^2(\mathbf{p}_1^2)}{64m_1^2m_2^2} \\ + \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{4m_1^2m_2^2} - \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_2)^2(\mathbf{p}_1 \cdot \mathbf{p}_2)^2}{16m_1^2m_2^2} - \frac{5(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{64m_1^2m_2^2} + \frac{21(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{64m_1^2m_2^2} \\ + \frac{3(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{32m_1^2m_2^2} - \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)^2(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1 \cdot \mathbf{p}_2)(\mathbf{p}_2^2)}{4m_1^2m_2^2} + \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{16m_1^2m_2^2} + \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1 \cdot \mathbf{p}_2)(\mathbf{p}_2^2)}{16m_1^2m_2^2} + \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1 \cdot \mathbf{p}_2)^2}{6m_1^2m_2^2} \\ + \frac{\mathbf{p}_1^2(\mathbf{p}_1 \cdot \mathbf{p}_2)(\mathbf{p}_2^2)}{32m_1^2m_2^2} + \frac{7(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{64m_1^2m_2^2} - \frac{3(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{32m_1^2m_2^2} - \frac{7(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{128m_1^2m_2^2}. \quad (\text{A4a})$$

$$H_{40}(\mathbf{x}_a, \mathbf{p}_a) = \frac{369(\mathbf{n}_{12} \cdot \mathbf{p}_1)^4}{160m_1^4} - \frac{889(\mathbf{n}_{12} \cdot \mathbf{p}_1)^2(\mathbf{p}_1^2)}{157m_1^4} + \frac{49(\mathbf{n}_{12} \cdot \mathbf{p}_1)^2(\mathbf{p}_1^2)^2}{16m_1^4} - \frac{63(\mathbf{p}_1^2)^3}{64m_1^4} - \frac{545(\mathbf{n}_{12} \cdot \mathbf{p}_1)^2(\mathbf{n}_{12} \cdot \mathbf{p}_2)}{128m_1^3m_2} \\ + \frac{67(\mathbf{n}_{12} \cdot \mathbf{p}_1)^2(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)}{15m_1^3m_2} - \frac{167(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)^2}{128m_1^2m_2} + \frac{1547(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1 \cdot \mathbf{p}_2)}{256m_1^2m_2} - \frac{831(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1 \cdot \mathbf{p}_2)}{128m_1^2m_2} \\ + \frac{1099(\mathbf{p}_1^2)^2(\mathbf{p}_1 \cdot \mathbf{p}_2)}{256m_1^2m_2} - \frac{3253(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)^2}{1280m_1^2m_2} + \frac{1067(\mathbf{n}_{12} \cdot \mathbf{p}_1)^2(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)}{480m_1^2m_2} - \frac{4567(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)^2}{3840m_1^2m_2} \\ + \frac{3571(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1 \cdot \mathbf{p}_2)}{350m_1^2m_2} + \frac{2073(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)(\mathbf{p}_1 \cdot \mathbf{p}_2)}{480m_1^2m_2} + \frac{4345(\mathbf{n}_{12} \cdot \mathbf{p}_1)^2(\mathbf{p}_1 \cdot \mathbf{p}_2)^2}{1280m_1^2m_2} \\ + \frac{3461(\mathbf{p}_1^2)(\mathbf{p}_1 \cdot \mathbf{p}_2)^2}{3840m_1^2m_2} + \frac{1673(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{p}_1^2)}{1280m_1^2m_2} - \frac{1939(\mathbf{n}_{12} \cdot \mathbf{p}_1)^2(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{3840m_1^2m_2} + \frac{2081(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{3840m_1^2m_2} - \frac{13(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)^2}{8m_1^2m_2} \\ + \frac{191(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)}{192m_1^2m_2} - \frac{19(\mathbf{n}_{12} \cdot \mathbf{p}_1)^2(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1 \cdot \mathbf{p}_2)}{384m_1^2m_2} + \frac{5(\mathbf{n}_{12} \cdot \mathbf{p}_2)^2(\mathbf{p}_1 \cdot \mathbf{p}_2)}{384m_1^2m_2} \\ + \frac{11(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1 \cdot \mathbf{p}_2)^2}{192m_1^2m_2} - \frac{77(\mathbf{p}_1 \cdot \mathbf{p}_2)^2}{96m_1^2m_2} + \frac{233(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{96m_1^2m_2} - \frac{47(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{32m_1^2m_2} \\ + \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{p}_1 \cdot \mathbf{p}_2)(\mathbf{p}_2^2)}{384m_1^2m_2} - \frac{18(\mathbf{p}_1^2)(\mathbf{p}_1 \cdot \mathbf{p}_2)(\mathbf{p}_2^2)}{384m_1^2m_2} - \frac{7(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{4m_1^2m_2} + \frac{7(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{4m_1^2m_2} \\ + \frac{7(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1 \cdot \mathbf{p}_2)}{2m_1^2m_2} - \frac{21(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1 \cdot \mathbf{p}_2)^2}{16m_1^2m_2} - \frac{7(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{6m_1^2m_2} - \frac{49(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{48m_1^2m_2} \\ + \frac{132(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1 \cdot \mathbf{p}_2)(\mathbf{p}_2^2)}{24m_1^2m_2} - \frac{77(\mathbf{p}_1 \cdot \mathbf{p}_2)(\mathbf{p}_2^2)}{96m_1^2m_2} + \frac{197(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{96m_1^2m_2} - \frac{173(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{48m_1^2m_2} + \frac{13(\mathbf{p}_1^2)^2}{8m_1^2}. \quad (\text{A4b})$$

$$H_{40}(\mathbf{x}_a, \mathbf{p}_a) = \frac{3127(\mathbf{n}_{12} \cdot \mathbf{p}_1)^4}{384m_1^4} - \frac{2295.3(\mathbf{n}_{12} \cdot \mathbf{p}_1)^2(\mathbf{p}_1^2)}{950m_1^4} - \frac{6695(\mathbf{p}_1^2)^2}{1152m_1^4} - \frac{3191(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1 \cdot \mathbf{p}_2)}{640m_1^3m_2} \\ + \frac{28561(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)}{1920m_1^3m_2} + \frac{8777(\mathbf{n}_{12} \cdot \mathbf{p}_1)^2(\mathbf{p}_1 \cdot \mathbf{p}_2)}{384m_1^3m_2} - \frac{752464(\mathbf{p}_1^2)(\mathbf{p}_1 \cdot \mathbf{p}_2)}{28800m_1^2m_2} \\ - \frac{16481(\mathbf{n}_{12} \cdot \mathbf{p}_1)^2(\mathbf{n}_{12} \cdot \mathbf{p}_2)^2}{960m_1^2m_2} + \frac{94433(\mathbf{n}_{12} \cdot \mathbf{p}_2)^2(\mathbf{p}_1^2)}{4300m_1^2m_2} - \frac{103957(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1 \cdot \mathbf{p}_2)}{2400m_1^2m_2} \\ + \frac{791(\mathbf{p}_1 \cdot \mathbf{p}_2)^2}{400m_1^2m_2} - \frac{26627(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{1600m_1^2m_2} - \frac{118261(\mathbf{p}_1^2)(\mathbf{p}_2^2)}{4800m_1^2m_2} - \frac{105(\mathbf{p}_1^2)^2}{32m_1^2}. \quad (\text{A4c})$$

$$H_{42}(\mathbf{x}_a, \mathbf{p}_a) = \left( \frac{2749\pi^2}{8.92} - \frac{211189}{19200} \right) \frac{(\mathbf{p}_1^2)^2}{m_1^4} + \left( \frac{63347}{1600} - \frac{1059\pi^2}{1024} \right) \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{p}_1^2)}{m^4} + \left( \frac{375\pi^2}{8192} - \frac{23533}{1280} \right) \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)^2}{m^4} \\ + \left( \frac{10631\pi^2}{8192} - \frac{1915349}{57600} \right) \frac{(\mathbf{p}_1 \cdot \mathbf{p}_2)^2}{m_1^2m_2^2} + \left( \frac{12723\pi^2}{16384} - \frac{2492417}{57600} \right) \frac{\mathbf{p}_1^2 \mathbf{p}_2^2}{m_1^2m_2^2} \\ + \left( \frac{1411429}{19200} - \frac{10092\pi^2}{512} \right) \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)}{m_1^2m_2^2} + \left( \frac{248991}{6400} - \frac{6153\pi^2}{2048} \right) \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1 \cdot \mathbf{p}_2)}{m_1^2m_2^2} \\ - \left( \frac{30383}{960} + \frac{36405\pi^2}{16384} \right) \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1 \cdot \mathbf{p}_2)^2}{m_1^2m_2^2} + \left( \frac{1243717}{14400} - \frac{40483\pi^2}{16384} \right) \frac{\mathbf{p}_1^2(\mathbf{p}_1 \cdot \mathbf{p}_2)}{m_1^2m_2^2} \\ + \left( \frac{2469}{60} + \frac{55655\pi^2}{16384} \right) \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)}{m_1^2m_2} + \left( \frac{4310\pi^2}{16384} - \frac{39111}{6400} \right) \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)(\mathbf{p}_1^2)}{m_1^2m_2} \\ + \left( \frac{56955\pi^2}{16384} - \frac{1645983}{12200} \right) \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)^2(\mathbf{p}_1 \cdot \mathbf{p}_2)}{m_1^2m_2}. \quad (\text{A4d})$$

$$H_{21}(\mathbf{x}_a, \mathbf{p}_a) = \frac{64861(\mathbf{p}_1^2)}{4800m_1^2} - \frac{91(\mathbf{p}_1 \cdot \mathbf{p}_2)}{8m_1m_2} + \frac{105(\mathbf{p}_2^2)}{32m_2^2} - \frac{9841(\mathbf{n}_{12} \cdot \mathbf{p}_1)^2}{1600m_1^2} - \frac{7(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)}{2m_1m_2}. \quad (\text{A4e})$$

$$H_{422}(\mathbf{x}_a, \mathbf{p}_a) = \left( \frac{1237033}{57600} - \frac{199177\pi^2}{49152} \right) \frac{\mathbf{p}_1^2}{m_1^4} + \left( \frac{176033\pi^2}{24576} - \frac{2864917}{57600} \right) \frac{(\mathbf{p}_1 \cdot \mathbf{p}_2)}{m_1m_2} + \left( \frac{282351}{19200} - \frac{21837\pi^2}{8152} \right) \frac{\mathbf{p}_2^2}{m_2^4} \\ + \left( \frac{698723}{19200} + \frac{21745\pi^2}{16384} \right) \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)^2}{m_1^4} - \left( \frac{63641\pi^2}{24576} - \frac{2712013}{19200} \right) \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)(\mathbf{n}_{12} \cdot \mathbf{p}_2)}{m_1m_2} \\ + \left( \frac{3200179}{57600} - \frac{28621\pi^2}{24576} \right) \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_2)^2}{m_2^4}. \quad (\text{A4f})$$

$$H_{43}(\mathbf{x}_a, \mathbf{p}_a) = -\frac{m_1^4}{16} + \left( \frac{6237\pi^2}{1024} - \frac{169799}{2400} \right) m_1^2 m_2 + \left( \frac{4825\pi^2}{6144} - \frac{603427}{7200} \right) m_1^2 m_2^2. \quad (\text{A4g})$$

$$H_{4PN}^{\text{nonloc}}(t) = -\frac{1}{5} \frac{G^2 M}{c^8} I_{ij}^{(3)}(t) \\ \times \text{Pf}_{2r_{12}/c} \int_{-\infty}^{+\infty} \frac{dv}{|v|} I_{ij}^{(3)}(t+v),$$

# Resummed (non-spinning) 4PN EOB interaction potentials

$$M = m_1 + m_2, \quad \mu = \frac{m_1 m_2}{m_1 + m_2}, \quad \nu = \frac{m_1 m_2}{(m_1 + m_2)^2} = \frac{\mu}{M} \quad u \equiv \frac{GM}{R c^2}$$

$$ds_{\text{eff}}^2 = -A(r; \nu) dt^2 + B(r; \nu) dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\varphi^2) \quad \bar{D} \equiv (A B)^{-1}$$

$$A(u) = 1 - 2u + 2\nu u^3 + \left( \frac{94}{3} - \frac{41\pi^2}{32} \right) \nu u^4 + \left( \left( \frac{2275\pi^2}{512} - \frac{4237}{60} + \frac{128}{5} \gamma_E + \frac{256}{5} \ln 2 \right) \nu + \left( \frac{41\pi^2}{32} - \frac{221}{6} \right) \nu^2 + \frac{64}{5} \nu \ln u \right) u^5,$$

$$A^{\text{EOB}}(u) = \text{Pade}_4^1[A^{\text{PN}}(u)]$$


$$\bar{D}(u) = 1 + 6\nu u^2 + (52\nu - 6\nu^2)u^3 + \left( \left( -\frac{533}{45} - \frac{23761\pi^2}{1536} + \frac{1184}{15} \gamma_E - \frac{6496}{15} \ln 2 + \frac{2916}{5} \ln 3 \right) \nu + \left( \frac{123\pi^2}{16} - 260 \right) \nu^2 + \frac{592}{15} \nu \ln u \right) u^4,$$

$$\hat{Q}(\mathbf{r}', \mathbf{p}') = \left( 2(4 - 3\nu)\nu u^2 + \left( \left( -\frac{5308}{15} + \frac{496256}{45} \ln 2 - \frac{33048}{5} \ln 3 \right) \nu - 83\nu^2 + 10\nu^3 \right) u^3 \right) (\mathbf{n}' \cdot \mathbf{p}')^4 + \left( \left( -\frac{827}{3} - \frac{2358912}{25} \ln 2 + \frac{1399437}{50} \ln 3 + \frac{390625}{18} \ln 5 \right) \nu - \frac{27}{5} \nu^2 + 6\nu^3 \right) u^2 (\mathbf{n}' \cdot \mathbf{p}')^6 + \mathcal{O}[\nu u (\mathbf{n}' \cdot \mathbf{p}')^8].$$

# Padé-resumming $A(u; \nu)$

$$u = \frac{GM}{c^2 r}$$

$$\nu = \frac{m_1 m_2}{(m_1 + m_2)^2}$$



# Resummed EOB waveform

(Damour-Iyer-Sathyaprakash '98) Damour-Nagar '07, Damour-Iyer -Nagar '08, Pan et al. '10

$$h_{\ell m} \equiv h_{\ell m}^{(N, \epsilon)} \hat{h}_{\ell m}^{(\epsilon)} \hat{h}_{\ell m}^{\text{NQC}}$$

$$\hat{h}_{\ell m}^{(\epsilon)} = \hat{S}_{\text{eff}}^{(\epsilon)} T_{\ell m} e^{i\delta_{\ell m}} \rho_{\ell m}^{\ell}$$

$$T_{\ell m} = \frac{\Gamma(\ell + 1 - 2i\hat{k})}{\Gamma(\ell + 1)} e^{\pi\hat{k}} e^{2i\hat{k} \ln(2kr_0)}$$

NB:  $T_{\ell m}$  resums an infinite number of terms and already contains, eg, 4.5PN tail<sup>3</sup> terms  
(Messina-Nagar17)

$$\begin{aligned} \rho_{22}(x; \nu) = & 1 + \left( \frac{55\nu}{84} - \frac{43}{42} \right) x + \left( \frac{19583\nu^2}{42336} - \frac{33025\nu}{21168} - \frac{20555}{10584} \right) x^2 \\ & + \left( \frac{10620745\nu^3}{39118464} - \frac{6292061\nu^2}{3259872} + \frac{41\pi^2\nu}{192} - \frac{48993925\nu}{9779616} - \frac{428}{105} \text{eulerlog}_2(x) + \frac{1556919113}{122245200} \right) x^3 \\ & + \left( \frac{9202}{2205} \text{eulerlog}_2(x) - \frac{387216563023}{160190110080} \right) x^4 + \left( \frac{439877}{55566} \text{eulerlog}_2(x) - \frac{16094530514677}{533967033600} \right) x^5 + \mathcal{O}(x^6), \end{aligned}$$

$$\mathcal{F}_{\varphi} \equiv -\frac{1}{8\pi\Omega} \sum_{\ell=2}^{\ell_{\text{max}}} \sum_{m=1}^{\ell} (m\Omega)^2 |Rh_{\ell m}^{(\epsilon)}|^2$$


# HAMILTON'S EQUATIONS & RADIATION REACTION

$$\dot{r} = \left(\frac{A}{B}\right)^{1/2} \frac{\partial \hat{H}_{\text{EOB}}}{\partial p_{r_*}}$$

$$\dot{\varphi} = \frac{\partial \hat{H}_{\text{EOB}}}{\partial p_{\varphi}} \equiv \Omega$$

$$\dot{p}_{r_*} = - \left(\frac{A}{B}\right)^{1/2} \frac{\partial \hat{H}_{\text{EOB}}}{\partial r} + \hat{\mathcal{F}}_{r_*}$$


$$\dot{p}_{\varphi} = \hat{\mathcal{F}}_{\varphi}$$



resummed conservative dynamics

resummed radiation reaction

# STRUCTURE OF THE EOB FORMALISM




$$h_{\ell m}^{\text{EOB}} = \theta(t_m - t) h_{\ell m}^{\text{insplunge}}(t) + \theta(t - t_m) h_{\ell m}^{\text{ringdown}}(t)$$

# Black Hole Quasi-Normal Modes (QNMs) (Vishveshwara'70, Press'71)


Perturbations of Schwarzschild spacetime: Regge-Wheeler57-Zerilli70 equation

$$\left[ \frac{\partial^2}{\partial r_*^2} - \frac{\partial^2}{\partial t^2} - V_\ell(r) \right] u_\ell(r, t) = 0. \quad r_* = r + 2M \log \left( \frac{r}{2M} - 1 \right)$$

$$V_\ell(r) = \left( 1 - \frac{2M}{r} \right) \left[ \frac{\ell(\ell+1)}{r^2} + \frac{2M}{r^3} \right]$$



**Quasi-Normal Modes: no incoming wave:  
linked to unstable light ring null geodesic at  $r_{LR} = 3 GM/c^2$   
complex frequency:  $\omega = \omega_0 - i \alpha$**



$$\sigma_N^+ = \alpha_N + i \omega_N$$

$$\left( \frac{Rc^2}{GM} \right) h_{22}^{\text{ringdown}}(t) = \sum_N C_N^+ e^{-\sigma_N^+(t-t_m)}$$